

§ 30.51

case of metal packages the tare established at the time of filling may be used unless it appears to be incorrect. From the proofs and the net weights of the packages, the wine gallons (if desired) and the proof gallons of spirits shall be determined by the use of Table 2. However, if the spirits contain solids in excess of 600 milligrams per 100 milliliters, the wine gallon and proof gallon contents shall be determined as prescribed for such spirits in § 30.41. If either the weight or the proof is beyond the limitations of table 2, either table 3 or table 4 may be used.

(Sec. 201, Pub. L. 85-859, 72 Stat. 1358, as amended (26 U.S.C. 5204))

DETERMINATION OF QUANTITY BY VOLUME

§ 30.51 Procedures for measurement of bulk spirits.

Where the quantity of spirits (including denatured spirits) in bulk is to be determined by volume as authorized by this chapter, the measurement shall be made in tanks, by meters as provided in 27 CFR part 19, or by other devices or methods authorized by the appropriate TTB officer, or as otherwise provided in this chapter, or such measurement may be made in tank cars or tank trucks if calibration charts for such conveyances are provided and such charts have been accurately prepared, and certified as accurate, by engineers or other persons qualified to calibrate such conveyances. Volumetric measurements in tanks shall be made only in accurately calibrated tanks equipped with suitable measuring devices, whereby the actual contents can be correctly ascertained. If the temperature of spirits (including denatured spirits) is other than the standard of 60 degrees Fahrenheit, gallonage determined by volumetric measurements shall be corrected to the standard temperature by means of table 7. In the case of denatured spirits, the temperature-correction factor for the proof of the spirits used in denaturation will give sufficiently accurate results, except that the temperature-correction factor used for specially denatured spirits, Formula No. 18, should be that given in table 7 for 100 proof spirits. When the quantity of spirits, in

27 CFR Ch. I (4-1-07 Edition)

wine gallons, has been determined by volumetric measurement, the number of proof gallons shall be obtained by multiplying the wine gallons by the proof of the spirits as determined under § 30.31.

Example Gauge glass reading inches—88.
Wine gallons per inch—48.96.
Temperature °F—72.
Proof of spirits—86.8.
Temperature correction factor (Table 7)—0.995.
48.96 W.G.×88=4308.48 wine gallons.
4308.48 W.G.×0.995=4286.94 wine gallons.
4286.94 W.G.×0.868=3721.06392=3721.1 proof gallons.

(Sec. 201, Pub. L. 85-859, 72 Stat. 1358, as amended (26 U.S.C. 5204))

[T.D. ATF-198, 50 FR 8535, Mar. 1, 1985, as amended by T.D. ATF-381, 61 FR 37004, July 16, 1996]

§ 30.52 Procedure for measurement of cased spirits.

Where the quantity of spirits in a case is to be determined by volume, such determination shall be made by ascertaining the contents of one bottle in the case and multiplying that figure by the number of bottles in the case. For cases containing bottles filled according to the metric system of measure, the quantity determined shall be converted to wine gallons, as provided in § 19.722 of this chapter. The wine gallons of spirits thus determined for one case may then be multiplied by the number of cases containing spirits at the same proof when determining the quantity of spirits for more than one case. The proof gallons of spirits in cases shall be determined by multiplying the wine gallons by the proof (divided by 100).

(Sec. 201, Pub. L. 85-859, 72 Stat. 1358, as amended (26 U.S.C. 5204))

Subpart E—Prescribed Tables

NOTE. The tables referred to in this subpart appear in their entirety in the “Gauging Manual Embracing Instructions and Tables for Determining Quantity of Distilled Spirits by Proof and Weight” which is incorporated by reference in this part (see § 30.1).